

The Initial Objects, Terminal Objects, Equalizers and Intersections on Categories of F_R^Λ -modules.

Zhao Jianli

Department of Mathematics, Liaocheng Teachers College, Shandong, 252059, P. R. China

Abstract: In this paper, we show that the category of F_R^Λ -modules is a top category, we obtain that the category of F_R^Λ -module has initial objects, terminal objects, equalizers and intersections.

Keywords: F_R^Λ -module, F -homomorphism, category of F_R^Λ -module, initial object, terminal object, top category, equalizer, intersection.

1 Introduction

[1], [2], [3] introduce the concepts of F_R^Λ -module and the category of F_R^Λ -modules, and discuss the properties of them. In this paper, we show that the category of F_R^Λ -modules is a top category, and we obtain that the category of F_R^Λ -module has initial objects, terminal objects, equalizers and intersections.

Let X be a nonempty set, L be a complete distributive lattice (with 0 and 1), A a fuzzy subset on X is characterised by a mapping $A: X \rightarrow L$. X^L denotes the set of whole fuzzy subset of X . In this paper R is a ring with identity $1 \neq 0$ and module which involved is an unitary left R -module.

Definition 1.1. [1] Let M be a left R -module, A a fuzzy subring of R and $A(0) = 1$, $B_M \in M^L$, if for all $x, y \in M, r \in R$, we have

$$1) B_M(x-y) \geq B_M(x) \wedge B_M(y),$$

$$2) B_M(0) = 1,$$

$$3) B_M(rx) \geq A(r) \wedge B(x),$$

then B_M is called an F_R^Λ -submodule (or F_R^Λ -module).

Definition 1.2. Let M and N be two R -modules, $f: M \rightarrow N$ be an R -homomorphism,

B_M be an F_R^\wedge -submodule of M , $\tilde{f}(B_M)$ is defined by

$$\tilde{f}(B_M)(y) = \begin{cases} \bigvee \{B_M(x) \mid x \in M, f(x) = y\}, & \text{if } f^{-1}(y) \neq \Phi, \\ 0, & \text{if } f^{-1}(y) = \Phi, \end{cases}$$

for all $y \in N$.

Definition 1. 3. Let M and N be two left R -module, $f: M \longrightarrow N$ be an R -homomorphism, B_M and C_N be F_R^\wedge -submodule of M and N , respectively, if $\tilde{f}(B_M) \leq C_N$, then \tilde{f} is called an F -homomorphism from B_M into C_N , writes $\tilde{f}: B_M \longrightarrow C_N$.

Definition 1. 4. The category of F_R^\wedge -modules $F_R^\wedge\text{-Mod}$ is defined by:

- 1) Objects are all F_R^\wedge -modules,
- 2) For all $B_M, C_N \in \text{Obj}(F_R^\wedge\text{-Mod})$, the set of morphisms is

$$\text{Hom}(B_M, C_N) = \{ \tilde{f} \mid \tilde{f} \text{ is an arbitrary } F\text{-homomorphism from } B_M \text{ into } C_N \},$$

- 3) For all $\tilde{f} \in \text{Hom}(B_M, C_N)$, $\tilde{g} \in \text{Hom}(C_N, D_S)$, the composition of \tilde{f} and \tilde{g} is defined by $\tilde{f}\tilde{g} = \tilde{fg}$.

Let $R\text{-Mod}$ denotes the category of left R -modules.

Definition 1. 5. Let $B_M, C_N \in \text{Obj}(F_R^\wedge\text{-mod})$, if $N \subseteq M$ and $B_M(x) \geq C_N(x)$, for all $x \in N$, then C_N is called the subobject of the object B_M .



2. Initial and terminal objects and top properties of F_R^\wedge -modules

Let M be a left R -module, A be a fuzzy subring of R , two F_R^\wedge -module of M are defined by

$$0_M: M \longrightarrow L, \quad 0_M(m) = \begin{cases} 0, & \text{if } m \neq 0, m \in M, \\ 1, & \text{if } m = 0, m \in M, \end{cases}$$

$$1_M: M \longrightarrow L, \quad 1_M(m) = 1, \forall m \in M.$$

Consequently, the diagram(1) and diagram(2) are admissible.



Hence the objects 0_M in category $F_R^\wedge\text{-Mod}$ is the initial objects of the category $F_R^\wedge\text{-Mod}$ and the objects $1_{\{0\}}$ is the terminal objects of the category $F_R^\wedge\text{-Mod}$, therefore we have the following Proposition 2. 1.

Proposition 2. 1. The category $F_R^\wedge\text{-Mod}$ has initial objects and terminal objects.

By Proposition 2. 1, we have the following Proposition 2. 2.

Proposition 2. 2. The category $F_R^\wedge\text{-Mod}$ has zero objects.

Theorem 2. 3. The category $F_R^\wedge\text{-Mod}$ is additive category, but it is not abel category.

Proof. By [2], [3] and Proposition 2. 2, the category $F_R^\wedge\text{-Mod}$ has products, coproducts, kernels and cokernels and zero objects, hence the category $F_R^\wedge\text{-Mod}$ is additive category.

Let $\tilde{g}: C_N \longrightarrow B_M$ be a subobject of B_M , if \tilde{g} is normal, then $C_N = \tilde{f}^{-1}(B_M)$. Hence for $M \neq 0$, $\tilde{1}: 0_M \longrightarrow 1_M$ is a subobjects of 1_M which is not a Kernel, therefore $F_R^\wedge\text{-Mod}$ is not an abel category.

Proposition 2. 4. Let M be a left R -module, the set $\Omega_L(M) = \{B_M \mid B_M \text{ is an } F_R^\wedge\text{-module of } M\}$ is a complete lattice under the following order relation:

$$B_M \leq C_M \text{ iff } B_M(x) \leq C_M(x), \text{ for all } x \in M.$$

Proof. Let $\{B_M^i \mid i \in I\} \subseteq \Omega_L(M)$

$$(\bigwedge_{i \in I} B_M^i)(x) = \bigwedge_{i \in I} B_M^i(x),$$

$$\bigvee_{i \in I} B_M^i(x) = \bigwedge_{i \in I} \{C_M(x) \mid C_M \in \Omega_L(M), B_M^i \leq C_M, \text{ for all } i \in I\},$$

are the inf and sub of the collection $\{B_M^i \mid i \in I\}$, respectively.

Theorem 2. 5. The category $F_R^\wedge\text{-Mod}$ is a top category over $R\text{-Mod}$.

Proof. The proof is similar to Theorem 3. 4 of [4].

3 Equalizers and intersections

Theorem 3. 1 The category $F_R^\wedge\text{-Mod}$ has equalizers.

Proof. Let $B_M, C_N \in \text{Obj}(F_R^\wedge\text{-Mod})$, and $\tilde{f}_1, \tilde{f}_2: B_M \longrightarrow C_N$ be two morphisms in $F_R^\wedge\text{-Mod}$. So we have two R -homomorphisms $f_1, f_2: M \longrightarrow N$ in $R\text{-Mod}$. But the category $R\text{-Mod}$ has equalizers and let the equalizer of f, g in $R\text{-Mod}$ be $i_k: K \longrightarrow M$, where

$$K = \{x \mid f_1(x) = f_2(x), x \in M\},$$

and i_K is the inclusion map. Evidently, K is an R -submodule of M . We define an F_R^A -module of K ,

$$D_K : K \longrightarrow L, D_K(x) = B_M(x), \forall x \in K,$$

since

$$D_K(x) \leq B_M(i_K(x)), \text{ for all } x \in K,$$

consequently, $\tilde{i}_K : D_K \longrightarrow B_M$ is an F -homomorphism. By the above construction we get that the following diagram holds in F_R^A -Mod.

$$\begin{array}{ccccc} D_K & \xrightarrow{\tilde{i}_K} & B_M & \xrightarrow{\tilde{f}_1} & H_N \\ & & & & \parallel \\ H_N & \xleftarrow{\tilde{f}_2} & B_M & \xleftarrow{\tilde{i}_K} & D_K \end{array}$$


Let there is an F -homomorphism $g : C'_K \longrightarrow B_M$ such that the following diagram holds in F_R^A -Mod.

$$\begin{array}{ccccc} D'_K & \xrightarrow{\tilde{g}} & B_M & \xrightarrow{\tilde{f}_1} & C_N \\ & & & & \parallel \\ C_N & \xleftarrow{\tilde{f}_2} & B_M & \xleftarrow{\tilde{g}} & D'_K \end{array}$$

So

$$\begin{array}{ccccc} K' & \xrightarrow{g} & M & \xrightarrow{f_1} & N \\ & & & & \parallel \\ N & \xleftarrow{f_2} & M & \xleftarrow{g} & K' \end{array}$$

holds in R -Mod. Since category R -Mod has equalizers, from the universal property there exists a unique R -homomorphism $k' : K' \longrightarrow K$ in R -Mod such that $i_K k' = g$. Since we have the following diagram(3)



is commutative. Therefore

$$D'_k(x) \leq B_M(x) \leq B_M(i_k k(x)) \\ = B_M i_k(k(x)) = D_k(x),$$

then \tilde{k}' is an F -homomorphism. Consequently, the following diagram holds

$$\begin{array}{ccc} D'_k & \xrightarrow{\tilde{k}} & D_k \xrightarrow{\tilde{i}_k} B_M \\ & & \parallel \\ & & B_M \xleftarrow{\tilde{g}} D'_k \end{array}$$

Hence the pair (D_k, \tilde{i}_k) is the equalizer of the pair of F -homomorphisms \tilde{f}_1 and \tilde{f}_2 in category $F_R^A\text{-Mod}$.

Theorem 3. 2. The category $F_R^A\text{-Mod}$ has finite intersections.

Proof. Let $\{B_{M_i}^i \mid i \in 1, 2, \dots, n\}$ be the family of subobjects of the object B_M in $F_R^A\text{-Mod}$.

Let $M' = \bigcap_{i=1}^n M_i$, we define fuzzy subset $B_{M'}$ of M' ,

$$B_{M'}: M' \longrightarrow L$$

such that

$$B_{M'}(x) = \bigwedge \{B_{M_i}^i(x) \mid i=1, \dots, n\}, \text{ for all } x \in M',$$

it is easy to prove $B_{M'} \in \text{Obj}(F_R^A\text{-Mod})$.

For all $i, g_i: M_i \longrightarrow M$ are inclusion map, because

$$B_{M_i}^i(x) \leq B_M(x)(f_i(x)), \text{ for any } x \in M_i, i=1, \dots, n$$

so for all $i=1, 2, \dots, n, \tilde{g}_i: B_{M_i}^i \longrightarrow B_M$ are F -homomorphism, let $\tilde{f}: D_H \longrightarrow B_M$ be an F -homomorphism which is factored through each subobject $B_{M_i}^i$, that is, for all $i \in I$, the following diagram holds.

$$\begin{array}{ccc} D_H & \xrightarrow{\tilde{f}_i} & B_{M_i}^i \xrightarrow{\tilde{g}_i} B_M \\ & & \parallel \\ & & B_M \xleftarrow{\tilde{f}} D_H \end{array}$$


Hence

$$\begin{array}{ccc} H & \xrightarrow{f_i} & M_i \xrightarrow{g_i} M \\ & & \parallel \\ & & M \xleftarrow{f} H \end{array}$$

holds, for $i=1, 2, \dots, n$. By the universal properties of intersection, there exists R -homomorphism $f': H \longrightarrow M'$ in $R\text{-Mod}$ such that

$$\begin{array}{ccccc} H & \xrightarrow{f'} & M' & \xrightarrow{g'} & M \\ & & & & \parallel \\ & & M & \xleftarrow{f} & H \end{array}$$

holds in $R\text{-Mod}$. Consider the following diagram(4)



For all $x \in M$, we have

$$\begin{aligned} D_H(x) &\leq B_{M_i}(f_i(x)) & \forall i=1,2,\dots,n, \\ g'_i f'_i &= f_i, g_i g'_i = g^i & \forall i=1,2,\dots,n, \end{aligned}$$

and $B_{M'}(x) = \bigwedge \{B_{M_i}(x) \mid i=1,2,\dots,n\}$. Then $D_H(x) \leq B_{M'}(f'(x))$, for all $x \in H$, so $\tilde{f}' : D_H \rightarrow B_{M'}$ is an F -homomorphism and

$$\begin{array}{ccccc} D_H & \xrightarrow{\tilde{f}'} & B_{M'} & \xrightarrow{\tilde{g}'} & B_M \\ & & & & \parallel \\ & & B_M & \xleftarrow{\tilde{f}} & D_H \end{array}$$

holds in $F_R^\wedge\text{-Mod}$. Therefore subobject $B_{M'}$ together with the family of F -homomorphisms

$$\{\tilde{g}'_i = B_{M'} \rightarrow B_{M_i} \mid i=1,2,\dots,n\}$$

is the intersection of the family of subobjects $\{B_{M_i} \mid i=1,\dots,n\}$ in $F_R^\wedge\text{-Mod}$.

Reference

[1] Zhao Jianli, Shi Kaiquan, Yu Mingshan, Fuzzy Modules over Fuzzy Rings, The J. Fuzzy Math, 3(1993), 531-539.
 [2] Zhao Jianli, F_R^\wedge -modules and F_R^\wedge -modules Categories The J. Fuzzy Math(to appear).
 [3] Zhao Jianli, Fuzzy homological theory in F_R^\wedge -modules(I), The J. Fuzzy Math (to appear).

- [4] S. R. LOPEZ—PERMOUTH and D. S. MALIK, On Categories of Fuzzy Modules, Inform Sci, (52)1990, 211—220.
- [5] S. M. A. Zaidi and Q. A. Ansari, Some results On categories of L—fuzzy subgroups of L—fuzzy subgroups, Fuzzy sets and systems 64(1994)249—256.
- [6] T. S. Blyth, Categories (Longman, London, 1987).
- [7] B. Mitchell, Theory of Categories (Academic Press, New York, 1965).