On fuzzy weakly semi-precontinuous multifunctions

Ma Baoguo

(Department of Mathematics and computer science, Yanan University, Yanan 716000 P.R.China)

Abstract: This paper is devoted to the introduction and study of fuzzy lower and upper weakly semi-precontinuous multifunctions between a general topological space (X,T) and a fuzzy topological space (Y,T_1) . Using the mapping F_* and F^* several equivelent conditions of a lower and upper fuzzy weakly semi-precontinuous multifunction.

Key words: Fuzzy semi-preopen (semi-preclosed) sets; Fuzzy lower and upper semi-precontinuous multifunctions; Fuzzy lower and upper weakly semi-precontinuous multifunctions.

1. Preliminaries

Throughout the paper , by (X,T) or simply by X we will mean a topological space in classical sense, and (Y,T_1) or simply by Y will stand for a fuzzy topological space (fts, for short) as defined by Chang's [1]. Fuzzy sets in Y will be denoted A,B,U,V etc. and interior(semi-interior) and closure (semi-closure) and complement of a fuzzy set A in an fts Y will be denoted by $Int\ A$ ($SInt\ A$) $Cl\ A$ ($SCl\ A$) and 1-A respectively. A fuzzy set A of fts X is called fuzzy preopen set (fuzzy preclosed set) if $A \le Int\ Cl\ A$ ($A \ge Int\ Cl\ A$). A fuzzy set A of fts X is called semi-preopen set (fuzzy semi-preclosed set) if there a fuzzy preopen set U such that $U \le A \le Cl\ U$ ($Int\ U \le A \le U$). Let A be any fuzzy set a fts X, then fuzzy semi-preclosure ($SPcl\ A$, for short) and fuzzy semi-preinterior (SP int A, for short) of A are defined as follows: $SPcl\ A = \bigcap \{B \mid B \text{ is fuzzy semi-preinterior} \text{ and } A \le B\}$ SP int $A = \bigcup \{B \mid B \text{ is fuzzy semi-preopen} \text{ and } B \le A\}$. when a fuzzy set A is quasi-coincident (q-coincident, for short) with a fuzzy set A in and $A \ge B$ are not q-coincident, denoted by

^{*} This work was partially supported by the special fundation of Shaanxi Educational committee.

 $A \neq B$. The words 'neighborhood' and 'fuzzy topological space' will be abbreviated as 'nbd', and 'fts' respectively.

Definition 1.1. Let (X,T) be a topological space in the classical sense and (Y,T_1) be a fuzzy topological space. $F:X\to Y$ is called a fuzzy multifunction iff for each $x\in X$, F(x) is a fuzzy set in Y.

Definition 1.2. For a fuzzy multifunction $F: X \to Y$, the $F_{\bullet}(A)$ and $F^{\bullet}(A)$ of a fuzzy set A in Y are defined as follows:

$$F_{\star}(A) = \{ x \in X : F(x) \neq A \},$$

$$F^{\star}(A) = \{ x \in X : F(x) \leq A \}$$

Theorem 1.3.[4] For a fuzzy multifunction $F: X \to Y$ we have $F_{\bullet}(1-A) = X - F^{\bullet}(A)$, for any fuzzy set A in Y.

Definition 1.4. For a fuzzy multifunction $F: X \to Y$ is called:

- (a) fuzzy lower semi-precontinuous (f.l.s.p.c., in short) at a point $x_0 \in X$ iff for every fuzzy open set V in Y will $x_0 \in F_{\bullet}(V)$, there exists a semi-preopen nbd U of x_0 in X such that $U \subset F_{\bullet}(V)$,
- (b) fuzzy upper semi-precontinuous (f.w.s.p.c., in short) at a point $x_0 \in X$ iff for every fuzzy open set V in Y will $x_0 \in F^*(V)$, there exists a semi-preopen nbd U of x_0 in X such that $U \subset F^*(V)$,
 - (c) f.l.s.p.c. (f.u.s.p.c.) on X iff it is respectively so at each $x_0 \in X$,
 - (d) f.s.p.c. on X iff it is f.l.s.p.c. and f.u.s.p.c.

Definition 1.5. For a fuzzy multifunction $F: X \to Y$ is called:

- (a) fuzzy lower weakly semi-continuous (f.l.w.s.c., in short) at a point $x_0 \in X$ iff for every fuzzy open set V in Y will $x_0 \in F_*(V)$, there exists a semi-open nbd U of x_0 in X such that $U \subset F_*(SCIV)$,
- (b) fuzzy upper weakly semi-continuous (f.u.w.s.c., in short) at a point $x_0 \in X$ iff for every fuzzy open set V in Y will $x_0 \in F^*(V)$, there exists a semi-open nbd U of x_0 in X such that $U \subset F^*(SCIV)$,
 - (c) f.l.w.s.c. (f.u.w.s.c.) on X iff it is respectively so at each $x_0 \in X$,
 - (d) f.w.s.c. on X iff it is f.l.w.s.c. and f.u.w.s.c.

2. Lower and upper weakly semi-precontinuous fuzzy multifunctions

Definition 2.1. For a fuzzy multifunction $F: X \to Y$ is called:

- (a) fuzzy lower weakly semi-precontinuous (f.l.w.s.p.c., in short) at a point $x_0 \in X$ iff for every fuzzy open set V in Y will $x_0 \in F$ (V), there exists a semi-preopen nbd U of x_0 in X such that $U \subset F$ (SCIV),
- (b) fuzzy upper weakly semi-precontinuous (f.u.w.s.p.c., in short) at a point $x_0 \in X$ iff for every fuzzy open set V in Y will $x_0 \in F^*(V)$, there exists a semi-preopen nbd U of x_0 in X such that $U \subset F^*(SCIV)$,
 - (c) f.l.w.s.p.c. (f.u.w.s.p.c.) on X iff it is respectively so at each $x_0 \in X$,
 - (d) f.w.s.p.c. on X iff it is f.l.w.s.p.c. and f.u.w.s.p.c.

Remark 2.2. It is clear from Difinition 2.1 and 1.4,1.5 that every fuzzy semi-continuous (weakly semi-continuous, almost semi-continuous [7], weakly semi-continuous [8], semi-precontinuous [9]) multifunctions is f.w.s.p.c. where as the converse is not true.

Theorem 2.3. Let $F: X \to Y$ is f.l.w.s.p.c.(f.u.w.s.p.c.) iff for every fuzzy open set V in Y, F. (V) \subset SP int F. (SCIV) (respectively, F* (V) \subset SP int F* (SCIV)).

Proof. we prove the theorem for the case of f.l.w.s.c. of F only; there other case is quite similar.

Let F be f.l.w.s.p.c. and V be f.open in Y. If $x \in F_*(V)$, then there exists semi-preopen nbd U of x in X such that $U \subseteq F_*(SCIV)$ and hence $U \subseteq SP$ int $F_*(SCIV)$.

Conversely, let $x \in X$ and V be fuzzy open set in Y such that $x \in F_{-}(V)$. Then $x \in F_{-}(V) \subset SP$ int $F_{-}(SCIV) = U$ (say). Thus U is a semi-preopen nbd of x such that $U \subset F_{-}(SCIV)$ and consequently, F is f.l.w.s.p.c..

Theorem 2.4. If $F: X \to Y$ is f.l.w.s.p.c., for every fuzzy pre-semiopen set V in $Y, F_*(V) \subset SP$ int $F_*(SCIV)$.

Proof. Let F be f.l.w.s.p.c. and V be any fuzzy pre-semiopen set in Y. For any $x \in F_*(V)$, then there exists semi-preopen nbd U_x of x in X such

that $U_x \subset F_*(SCIIntSCIV) \subset F_*(SCIV)$ and hence $F_*(V) \subset \bigcup \{U_x : x \in F_*(V)\} \subset F_*(SCIV)$, and hence $F^*(V) \subset SP$ int $F^*(SCIV)$.

Theorem 2.5. If $F: X \to Y$ is f.u.w.s.p.c., for every fuzzy pre-semiopen set V in Y, $F^*(V) \subset SP$ int $F^*(SCIV)$.

Proof. Similar to Theorem 2.4, and omitted.

Theorem 2.6. If a fuzzy multifunction $F: X \to Y$ is f.l.w.s.p.c. on X, then $SPcl\ F_{\bullet}(V) \subset F_{\bullet}(SClV)$ for any fuzzy open set V in Y.

Proof. Let $x \notin F_{\bullet}(SCIV)$, where V is a fuzzy open set in Y. Then F(x) = qSPcIV so $F(x) \le 1 - SPcIV$. By fuzzy lower weakly semi-precontinuity of F, there exists a semi-preopen nbd U of x in X such that $F(z) \le SPcI(1 - SCIV) \le 1 - V$ for all $z \in U$. Thus F(z) = qV, i.e. $z \notin F_{\bullet}(V)$, for all $z \in U$, hence $U = \bigcap F_{\bullet}(V) = \emptyset$. Since U is a semi-preopen nbd of x, it then follows that $x \notin F_{\bullet}(SCIV)$.

Theorem 2.7. If a fuzzy multifunction $F: X \to Y$ is f.u.w.s.p.c. on X, then $SPcl\ F^*(V) \subset F^*(SCIV)$ for any fuzzy open set V in Y.

Proof. Similar to Theorem 2.6. and omitted.

Theorem 2.8. Let $F: X \to Y$ is fuzzy multifunctions then following are equivalent:

- (a) F is f.l.w.s.p.c.,
- (b) for any fuzzy open set V in Y, $F_{\bullet}(V) \subset SP$ int $F_{\bullet}(SCIV)$ and $F^{*}(V) \subset F^{*}(SCIV)$.

Proof. Evident.

Theorem 2.9. Let $F: X \to Y$ is fuzzy lower weakly semi-precontinuity if for each fuzzy open set V, $F_{\star}(CIV)$ is fuzzy semi-preopen set.

Proof. Straightforward.

References

- [1] C.L.Chang, Fuzzy topological spaces, J.Math.Anal.Appl 24(1968) 182-190.
- [2] N.S.Papageorgiou, Fuzzy topology and fuzzy multifunction, J.Math.Anal. Appl. 109(1985) 397-425.
- [3] M.N.Mukherjee and S.Malaker, On almost continuous and weakly continuous fuzzy multifunctions, Fuzzy set and systems, 41(1991) 113-125.
- [4] B.G. Ma, On fuzzy semi-continuous and irresolute multifunction, © ICIK95 August 21-25, 1995, Dalian, P.R.China.
- [5] S.Dang, A.Behera. S.Nanda, On fuzzy weakly semi-continuous functions, Fuzzy set and systems, 67(1994) 239-245.
- [6] L.A.Zadeh, Fuzzy sets, Inform and Control 8(1965) 338-353.
- [7] B.G. Ma, Lower and upper almost semi-continuous fuzzy multifunction, Fuzzy Sets Theory and Applications (Hebei University Press 1998.8, Baoding P.R.China) 369-372.
- [8] B.G. Ma, Lower and upper wekly semi-continuous fuzzy multifunction, BUSEFAL(in France), Vol. 78(1999) 8-11.
- [9] B.G. Ma, Fuzzy semi-preopen sets and fuzzy semi-precontinuous mulitifunctions, Journal of yanan university (in chinese), 2(1999).